

Geometrically Uniform Frames

Yonina C. Eldar* and Helmut Bölcskei†

February 1, 2008

Abstract

We introduce a new class of frames with strong symmetry properties called *geometrically uniform frames (GU)*, that are defined over an abelian group of unitary matrices and are generated by a single generating vector. The notion of GU frames is then extended to *compound GU (CGU)* frames which are generated by an abelian group of unitary matrices using multiple generating vectors.

The dual frame vectors and canonical tight frame vectors associated with GU frames are shown to be GU and therefore generated by a single generating vector, which can be computed very efficiently using a Fourier transform defined over the generating group of the frame. Similarly, the dual frame vectors and canonical tight frame vectors associated with CGU frames are shown to be CGU.

The impact of removing single or multiple elements from a GU frame is considered. A systematic method for constructing optimal GU frames from a given set of frame vectors that are not GU is also developed. Finally, the Euclidean distance properties of GU frames are discussed and conditions are derived on the abelian group of unitary matrices to yield GU frames with strictly positive distance spectrum irrespective of the generating vector.

*Research Laboratory of Electronics, Massachusetts Institute of Technology, Room 36-615, Cambridge, MA 02139. E-mail: yonina@mit.edu.

†Coordinated Science Laboratory, University of Illinois at Urbana-Champaign, Urbana, IL 61801. Email: bolcskei@comm.csl.uiuc.edu.

1 Introduction

Frames are generalizations of bases which lead to redundant signal expansions [1, 2]. A (finite) frame for a Hilbert space $\mathcal{H}$ is a set of not necessarily linearly independent vectors that spans $\mathcal{H}$. Since the frame vectors can be linearly dependent, the conditions on frame vectors are usually not as stringent as the conditions on bases, allowing for increased flexibility in their design [3, 4].

Frames were first introduced by Duffin and Schaeffer [1] in the context of nonharmonic Fourier series, and play an important role in the theory of nonuniform sampling [1, 2, 5] and wavelet theory [3, 6]. Recently, frames have been used to analyze and design oversampled filter banks [7, 8, 9] and error correction codes [10]. Frames have also been applied to the development of modern uniform and nonuniform sampling techniques [11], to various detection problems [12, 13], and to multiple description source coding [14].

Two important classes of highly structured frames are Gabor (Weyl-Heisenberg (WH)) frames [15, 16] and wavelet frames [3, 6, 17]. Both classes of frames are generated by a single generating function. WH frames are obtained by translations and modulations of the generating function (referred to as the window function), and wavelet frames are obtained by shifts and dilations of the generating function (referred to as the mother wavelet). In Section 3 of this paper, we introduce a new class of frames which we refer to as geometrically uniform (GU) frames, that like WH and wavelet frames are generated from a single generating vector. These frames are defined by an abelian group $\mathcal{Q}$ of unitary matrices, referred to as the generating group of the frame. We note that WH frames and wavelet frames are in general not GU since the underlying group of matrices is in general not abelian. GU frames are based on the notion of GU vector sets first introduced by Slepian [18] and later extended by Forney [19], which are known to have strong symmetry properties that may be desirable in various applications such as channel coding [19, 20, 21].

The notion of GU frames is then extended to frames that are generated by an abelian group $\mathcal{Q}$ of unitary matrices using *multiple* generating vectors. Such frames are not necessarily GU, but

consist of subsets of GU vector sets that are each generated by $\mathcal{Q}$. We refer to this class of frames as compound GU (CGU) frames, and develop their properties in Section 6. CGU frames are a generalization of filter bank frames introduced in [7, 8, 9]. An interesting class of frames results when the set of generating vectors is itself GU, generated by an abelian group $\mathcal{G}$. (Note that this class of frames will in general not be GU). As we show, these frames are a generalization of WH frames in which $\mathcal{Q}$ is the group of translations and $\mathcal{G}$ is the group of modulations.

Given a frame for $\mathcal{H}$, any signal in $\mathcal{H}$ can be represented as a linear combination of the frame vectors. However, if the frame vectors are linearly dependent, then the coefficients in this expansion are not unique. A popular choice of coefficients are the inner products of the signal with a set of analysis frame vectors called the dual frame vectors [17]. This choice of coefficients has the property that among all possible coefficients it has the minimal l_2 -norm [17, 22].

In Section 4, we show that the dual frame vectors associated with a GU frame are also GU, and therefore generated by a single generating vector. Furthermore, we demonstrate that the generating vector can be computed very efficiently using a Fourier transform defined over the generating group $\mathcal{Q}$ of the frame. Similarly, in Section 6 we show that the dual frame vectors associated with a CGU frame are also CGU. When the generating vectors of the CGU frame are GU and generated by a group $\mathcal{G}$ that commutes up to a phase factor with the group $\mathcal{Q}$, the dual frame is generated by a *single* generating vector, a result well known for WH frames.

An important topic in frame theory is the behavior of a frame when elements of the frame are removed. In Section 7, we show that the frame bounds of the frame resulting from removing a single vector of a GU frame are the same regardless of the particular vector removed. In this sense GU frames exhibit an interesting robustness property which is of particular importance in applications such as multiple description source coding [14]. We also consider the behavior of a GU frame when groups of frame elements are removed.

In the special case of a tight frame the dual frame vectors are proportional to the original

frame vectors so that the reconstruction formula is particularly simple. In many applications it is therefore desirable to construct a tight frame from an arbitrary set of frame vectors. A popular tight frame construction is the so-called canonical tight frame [17, 7, 23, 24, 25, 26], first proposed in the context of wavelets in [27]. The canonical tight frame is relatively simple to construct, it is optimal in a least-squares sense [28, 26, 29], it can be determined directly from the given vectors, and plays an important role in wavelet theory [30, 31, 32]. Like the dual frame vectors, we show that the canonical tight frame vectors associated with a GU frame are GU, and the canonical tight frame vectors associated with a CGU frame are CGU. When the generating vectors of the CGU frame are GU and generated by a group $\mathcal{G}$ that commutes up to a phase factor with $\mathcal{Q}$, the canonical tight frame vectors can be obtained by a single generating vector, generalizing a result well-known in WH frame theory.

Since GU frames have nice symmetry properties, it may be desirable to construct such a frame from a given set of frame vectors. The problem of frame design has received relatively little attention in the literature. Systematic methods for constructing optimal tight frames have been considered [28, 26, 29]. Methods for generating frames starting from a given frame are described in [4]. In Section 8, we systematically construct optimal GU frames from a given set of vectors, that are closest in a least-squares sense to the original frame vectors. We consider three different constraints on the GU frame vectors. First, we treat the case in which the inner products of the frame vectors are known. The optimizing frame is referred to as the scaled-constrained least-squares GU frame (SC-LSGUF). Next, we consider the case where the inner products are known up to a scale factor. The optimizing frame in this case is referred to as the constrained least-squares GU frame (C-LSGUF). Finally, we consider the case in which both the inner products and the scaling are chosen to minimize the least-squares error between the original frame and the resulting tight frame. The optimizing frame is the least-squares GU frame (LSGUF).

In Section 9, we consider distance properties of GU frames, which may be of interest when using

GU frames for code design (group codes) [18, 19]. In particular, we introduce a class of GU frames with strictly positive distance spectra for all choices of generating vectors. Such GU frames are shown to be generated by fixed-point-free groups [33].

Before proceeding to the detailed development, in Section 2 we provide a brief introduction to frame expansions.

2 Frames

Frames, which are generalizations of bases, were introduced in the context of nonharmonic Fourier series by Duffin and Schaeffer [1] (see also [2]). Recently, the theory of frames has been expanded [3, 6, 17, 4], in part due to the utility of frames in analyzing wavelet decompositions.

Let $\{\phi_i, 1 \leq i \leq n\}$ denote a set of n complex vectors in an m -dimensional Hilbert space $\mathcal{H}$. The vectors ϕ_i form a *frame* for $\mathcal{H}$ if there exist constants $A > 0$ and $B < \infty$ such that

$$A\|x\|^2 \leq \sum_{i=1}^n |\langle x, \phi_i \rangle|^2 \leq B\|x\|^2, \quad (1)$$

for all $x \in \mathcal{H}$ [17]. In this paper, we restrict our attention to the case where m and n are finite. The lower bound in (1) ensures that the vectors ϕ_i span $\mathcal{H}$; thus we must have $n \geq m$. Since $n < \infty$ the right hand inequality of (1) is always satisfied with $B = \sum_{i=1}^n \|\phi_i\|^2$, so that any finite set of vectors that spans $\mathcal{H}$ is a frame for $\mathcal{H}$. In particular, any basis for $\mathcal{H}$ is a frame for $\mathcal{H}$. However, in contrast to basis vectors which are linearly independent, frame vectors with $n > m$ are linearly dependent. If the bounds $A = B$ in (1), then the frame is called a *tight frame*. If in addition $A = B = 1$, then the frame is called a *normalized tight frame*. The redundancy of the frame is defined as $r = n/m$, *i.e.*, n vectors in an m -dimensional space.

The *frame operator* corresponding to the frame vectors $\{\phi_i, 1 \leq i \leq n\}$ is defined as [17]

$$S = \sum_{i=1}^n \phi_i \phi_i^* = \Phi \Phi^*, \quad (2)$$

where Φ is the matrix of columns ϕ_i , and $(\cdot)^*$ denotes the Hermitian transpose. Using the frame operator, (1) can be rewritten as

$$A\|x\|^2 \leq \langle Sx, x \rangle \leq B\|x\|^2. \quad (3)$$

From (3) it follows that the tightest possible frame bounds A and B are given by $A = \min_i \lambda_i(S)$ and $B = \max_i \lambda_i(S)$, where $\{\lambda_i(S), 1 \leq i \leq m\}$ are the eigenvalues of the frame operator S . Throughout the paper, when referring to “the frame bounds” we implicitly assume the tightest possible frame bounds unless otherwise stated.

If the vectors $\{\phi_i, 1 \leq i \leq n\}$ form a frame for $\mathcal{H}$, then any $x \in \mathcal{H}$ can be expressed as a linear combination of these vectors: $x = \sum_{i=1}^n a_i \phi_i$. If $n > m$ the coefficients in this expansion are not unique. A possible choice is¹ $a_i = \langle \bar{\phi}_i, x \rangle$ where $\bar{\phi}_i$ are the *dual frame vectors* [17] of the frame vectors ϕ_i , and are given by

$$\bar{\phi}_i = S^{-1} \phi_i. \quad (4)$$

We can express S^{-1} directly in terms of S as [6]

$$S^{-1} = \frac{2}{A+B} \left[I - \left(I - \frac{2}{A+B} S \right) \right]^{-1} = \frac{2}{A+B} \sum_{l=0}^{\infty} \left(I - \frac{2}{A+B} S \right)^l, \quad (5)$$

where A and B are the frame bounds of $\{\phi_i\}$. The choice of coefficients $a_i = \langle \bar{\phi}_i, x \rangle$ has the property that among all possible coefficients it has the minimal l_2 -norm [17, 22].

There are other choices of dual frame vectors y_i such that for any $x \in \mathcal{H}$, $x = \sum_{i=1}^n \langle y_i, x \rangle \phi_i$.

¹We use the notation $\langle x, y \rangle = x^* y$.

Specifically, with Y denoting the matrix of columns y_i , any other choice corresponds to Y of the form [34]

$$Y = S^{-1}\Phi + T(I - \Phi^*S^{-1}\Phi), \quad (6)$$

where T is an arbitrary matrix with bounded elements. However, the particular choice $y_i = \bar{\phi}_i$ has some desirable properties. Besides resulting in the minimal l_2 -norm coefficients, in many cases the choice $y_i = \bar{\phi}_i$ yields frame vectors that share the same symmetries as the original frame vectors. Specifically, in Section 4 we show that the dual frame vectors associated with a geometrically uniform (GU) frame are also GU, and in Section 6 we show that the dual frame vectors associated with a compound GU (CGU) frame are also CGU. Finally, in the case of a tight frame the dual frame vectors lead to a particularly simple expansion. Specifically, in this case $S = AI_m$ so that $S^{-1} = (1/A)I_m$, and the dual frame vectors are $\{\bar{\phi}_i = (1/A)\phi_i, 1 \leq i \leq n\}$. Since a tight frame expansion of a signal is very simple, it is popular in many applications [17].

Suppose we are given a set of vectors $\{\phi_i, 1 \leq i \leq n\}$ that form a frame for $\mathcal{H}$, with frame bounds $A \neq B$. It may then be desirable to construct a tight frame from these vectors. A popular tight frame construction is the canonical tight frame [17, 7, 23, 24, 26, 28], first proposed in the context of wavelets in [27]. The *canonical tight frame vectors* $\{\mu_i, 1 \leq i \leq n\}$ associated with the vectors $\{\phi_i, 1 \leq i \leq n\}$ are given by

$$\mu_i = S^{-1/2}\phi_i, \quad (7)$$

where $S^{-1/2}$ is the positive-definite square root of S^{-1} . We can express $S^{-1/2}$ directly in terms of S as [35]

$$S^{-1/2} = \sqrt{\frac{2}{A+B}} \sum_{l=0}^{\infty} \frac{(2l)!}{2^{2l}(l!)^2} \left(I - \frac{2}{A+B}S\right)^l, \quad (8)$$

where A and B are the frame bounds associated with the frame $\{\phi_i\}$. Note, that $\{US^{-1/2}\phi_i\}$ with U an arbitrary unitary matrix yields a tight frame as well. The canonical tight frame, however, has the property that it is the closest normalized tight frame to the vectors $\{\phi_i\}$ in a least-squares

sense [25, 26, 29].

From (4) and (7) we see that in order to compute the dual frame vectors and the canonical tight frame vectors associated with a frame $\{\phi_i\}$, we need to compute the matrices S^{-1} and $S^{-1/2}$ and then apply them to each of the frame vectors ϕ_i . In the next section, we introduce a class of frames that have strong symmetry properties called *geometrically uniform (GU) frames*. As we show in Section 4, the dual frame vectors and the canonical tight frame vectors associated with a GU frame are generated by a single generating function, and can therefore be computed very efficiently.

3 Geometrically Uniform Frames

A set of vectors $\mathcal{S} = \{\phi_i, 1 \leq i \leq n\}$ is geometrically uniform (GU) [19, 18, 25] if every vector in the set has the form $\phi_i = U_i\phi$, where ϕ is an arbitrary *generating vector* and the matrices $\{U_i, 1 \leq i \leq n\}$ are unitary and form an abelian group² $\mathcal{Q}$. For concreteness we assume that $U_1 = I$ so that $\phi_1 = \phi$. The group $\mathcal{Q}$ will be called the *generating group* of $\mathcal{S}$.

Alternatively, a vector set is GU if given any two vectors ϕ_i and ϕ_j in the set, there is an isometry (a norm-preserving linear transformation) Z_{ij} that transforms ϕ_i into ϕ_j while leaving the set invariant [19]. Thus for every i , $Z_{ij}\phi_i = \phi_j$. Intuitively, a vector set is GU if it “looks the same” geometrically from any of the points in the set. Some examples of GU vector sets are considered in [19].

A set of vectors $\{\phi_i \in \mathcal{H}, 1 \leq i \leq n\}$ forms a *geometrically uniform frame* for $\mathcal{H}$, if the vectors $\{\phi_i\}$ are GU and span $\mathcal{H}$.

As we show in the following proposition, the frame bounds of a GU frame can be bounded by the norm of the generating vector.

Proposition 1. *Let $\mathcal{S} = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}\}$ be a geometrically uniform frame with frame bounds*

²That is, $\mathcal{Q}$ contains the identity matrix I ; if $\mathcal{Q}$ contains U_i , then it also contains its inverse U_i^{-1} ; the product $U_i U_j$ of any two elements of $\mathcal{Q}$ is again in $\mathcal{Q}$; and $U_i U_j = U_j U_i$ for any two elements in $\mathcal{Q}$ [36].

A and B , where ϕ is an arbitrary generating vector. Then $A \leq \frac{n}{m} \|\phi\|^2 \leq B$. If in addition the frame is tight, then $A = B = \frac{n}{m} \|\phi\|^2$.

Proof. We can express the frame operator corresponding to the frame vectors ϕ_i as

$$S = \sum_{i=1}^n U_i \phi \phi^* U_i^*. \quad (9)$$

Then,

$$\text{Tr}(S) = \sum_{i=1}^n \text{Tr}(U_i \phi \phi^* U_i^*) = n \|\phi\|^2, \quad (10)$$

so that

$$\sum_{i=1}^m \lambda_i(S) = \text{Tr}(S) = n \|\phi\|^2. \quad (11)$$

Therefore,

$$A = \min_i \lambda_i(S) \leq \frac{1}{m} \sum_{i=1}^m \lambda_i(S) = \frac{n}{m} \|\phi\|^2, \quad (12)$$

and

$$B = \max_i \lambda_i(S) \geq \frac{1}{m} \sum_{i=1}^m \lambda_i(S) = \frac{n}{m} \|\phi\|^2. \quad (13)$$

□

Since $U_i^* = U_i^{-1}$, the inner product of two vectors in $\mathcal{S}$ is

$$\langle \phi_i, \phi_j \rangle = \phi^* U_i^{-1} U_j \phi = s(U_i^{-1} U_j), \quad (14)$$

where s is the function on $\mathcal{Q}$ defined by

$$s(U_i) = \phi^* U_i \phi. \quad (15)$$

For fixed i , the set $U_i^{-1} \mathcal{Q} = \{U_i^{-1} U_j, U_j \in \mathcal{Q}\}$ is just a permutation of $\mathcal{Q}$ since $U_i^{-1} U_j \in \mathcal{Q}$ for

all i, j [36]. Therefore, the n numbers $\{s(U_i^{-1}U_j), 1 \leq j \leq n\}$ are a permutation of the numbers $\{s(U_i), 1 \leq i \leq n\}$. The same is true for fixed j . Consequently, every row and column of the $n \times n$ Gram matrix $G = \{\langle \phi_i, \phi_j \rangle\}$ is a permutation of the numbers $\{a_i = s(U_i), 1 \leq i \leq n\}$.

A matrix G whose rows (columns) are a permutation of the first row (column) will be called a *permuted matrix*³. Thus, we have shown that the Gram matrix of a GU vector set is a permuted matrix. Furthermore, if the Gram matrix $G = \{\langle \phi_i, \phi_j \rangle\}$ is a permuted matrix and in addition $G = G^T$, then the vectors $\{\phi_i\}$ are GU [37]. We therefore have the following proposition.

Proposition 2. *The Gram matrix $G = \{\langle \phi_i, \phi_j \rangle\}$ corresponding to a geometrically uniform vector set $\mathcal{S} = \{\phi_i \in \mathcal{H}, 1 \leq i \leq n\}$ is a permuted matrix. Conversely, if the Gram matrix $G = \{\langle \phi_i, \phi_j \rangle\}$ is a permuted matrix, and $\langle \phi_i, \phi_j \rangle = \langle \phi_j, \phi_i \rangle$ for all i, j , then the vectors $\{\phi_i\}$ are geometrically uniform. If in addition the vectors $\{\phi_i\}$ span $\mathcal{H}$, then they form a geometrically uniform frame for $\mathcal{H}$.*

It will be convenient to replace the multiplicative group $\mathcal{Q}$ by an additive group Q to which $\mathcal{Q}$ is isomorphic⁴. Specifically, it is well known (see *e.g.*, [36]) that every finite abelian group $\mathcal{Q}$ is isomorphic to a direct product Q of a finite number of cyclic groups: $\mathcal{Q} \cong Q = \mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_p}$, where $\mathbb{Z}_{n_t}$ is the cyclic additive group of integers modulo n_t , and $n = \prod_t n_t$. Thus every element $U_i \in \mathcal{Q}$ can be associated with an element $q \in Q$ of the form $q = (q_1, q_2, \dots, q_p)$, where $q_t \in \mathbb{Z}_{n_t}$; this correspondence is denoted by $U_i \leftrightarrow q$.

Each vector $\phi_i = U_i \phi$ is then denoted as $\phi(q)$, where $U_i \leftrightarrow q$. The zero element $0 = (0, 0, \dots, 0) \in Q$ corresponds to the identity matrix $I \in \mathcal{Q}$, and an additive inverse $-q \in Q$ corresponds to a

³An example of a permuted matrix is

$$\begin{bmatrix} a_1 & a_2 & a_3 & a_4 \\ a_2 & a_1 & a_4 & a_3 \\ a_3 & a_4 & a_1 & a_2 \\ a_4 & a_3 & a_2 & a_1 \end{bmatrix}. \quad (16)$$

⁴ Two groups $\mathcal{Q}$ and $\mathcal{Q}'$ are *isomorphic*, denoted by $\mathcal{Q} \cong \mathcal{Q}'$, if there is a bijection (one-to-one and onto map) $\varphi : \mathcal{Q} \rightarrow \mathcal{Q}'$ which satisfies $\varphi(xy) = \varphi(x)\varphi(y)$ for all $x, y \in \mathcal{Q}$ [36].

multiplicative inverse $U_i^{-1} = U_i^* \in \mathcal{Q}$. The Gram matrix is then the $n \times n$ matrix

$$G = \{\langle \phi(q'), \phi(q) \rangle, q', q \in Q\} = \{s(q - q'), q', q \in Q\}, \quad (17)$$

with row and column indices $q', q \in Q$, where s is now the function on Q defined by

$$s(q) = \langle \phi(0), \phi(q) \rangle. \quad (18)$$

The Fourier transform (FT) of a complex-valued function $\varphi : Q \rightarrow \mathbb{C}$ defined on $Q = \mathbb{Z}_{n_1} \times \cdots \times \mathbb{Z}_{n_p}$ is the complex-valued function $\hat{\varphi} : Q \rightarrow \mathbb{C}$ defined by

$$\hat{\varphi}(h) = \frac{1}{\sqrt{n}} \sum_{q \in Q} \langle h, q \rangle \varphi(q), \quad (19)$$

where the Fourier kernel $\langle h, q \rangle$ is

$$\langle h, q \rangle = \prod_{t=1}^p e^{-2\pi i h_t q_t / n_t}. \quad (20)$$

Here h_t and q_t are the k th components of h and q respectively, and the product $h_t q_t$ is taken as an ordinary integer modulo n_t .

The FT matrix over Q is defined as the $n \times n$ matrix $\mathcal{F} = \{\frac{1}{\sqrt{n}} \langle h, q \rangle, h, q \in Q\}$. The FT of a column vector $\varphi = \{\varphi(q), q \in Q\}$ is then the column vector $\hat{\varphi} = \{\hat{\varphi}(h), h \in Q\}$ given by $\hat{\varphi} = \mathcal{F}\varphi$. Since $\mathcal{F}$ is unitary, we obtain the inverse FT formula

$$\varphi = \mathcal{F}^* \hat{\varphi} = \left\{ \frac{1}{\sqrt{n}} \sum_{h \in Q} \langle h, q \rangle^* \hat{\varphi}(h), q \in Q \right\}. \quad (21)$$

As we show in the following theorem, the FT matrix plays an important role in defining GU frames.

Theorem 1. *A set of vectors $\{\phi_i, 1 \leq i \leq n\}$ in an m -dimensional Hilbert space $\mathcal{H}$ is geometrically uniform if and only if the Gram matrix $G = \{\langle \phi_i, \phi_j \rangle\}$ is diagonalized by a Fourier transform matrix $\mathcal{F}$ over a finite product of cyclic groups Q . The vectors $\{\phi_i\}$ form a geometrically uniform frame for $\mathcal{H}$ if in addition G has rank m .*

Proof. The vectors $\{\phi_i\}$ form a frame for $\mathcal{H}$ if and only if they span $\mathcal{H}$, which implies that the rank of G must be equal to m .

For a GU vector set with generating group $\mathcal{Q} \cong Q$, the FT over Q diagonalizes the Gram matrix G [25]. Thus, to complete the proof of the theorem we need to prove that if G is diagonalized by a FT matrix $\mathcal{F}$ over the group Q , then the vector set $\{\phi(q), q \in Q\}$ is GU.

Let Φ be the matrix of columns $\phi(q)$, so that $G = \Phi^* \Phi$. Since $\mathcal{F}$ diagonalizes G , G has an eigendecomposition of the form $G = \mathcal{F} D \mathcal{F}^*$ for a diagonal matrix D with diagonal elements d_i , where the first m diagonal elements may be non-zero and the remaining diagonal elements are all zero. Then Φ has an SVD of the form $\Phi = U \Sigma \mathcal{F}^*$ where U is an arbitrary unitary matrix and Σ is an $m \times n$ diagonal matrix with diagonal elements $\{\sigma_i = \sqrt{d_i}, 1 \leq i \leq m\}$.

Let $\{f(q), q \in Q\}$ denote the columns of $\mathcal{F}^*$. From the definition of $\mathcal{F}$, the components of $f(0)$ are all equal $1/\sqrt{n}$, and $f(q) = B(q)f(0)$ where $B(q)$ is a diagonal unitary matrix with diagonal elements $\{\langle h, q \rangle, h \in Q\}$, where $\langle h, q \rangle$ is given by (20). Then,

$$\phi(q) = U \Sigma f(q) = U \Sigma B(q) f(0) = U B(q) \Sigma f(0) = U B(q) U^* \phi, \quad (22)$$

where $\phi = U \Sigma f(0)$, and where we used the fact that diagonal matrices commute. If we now define $U(q) = U B(q) U^*$, then we have that $\phi(q) = U(q) \phi$ where the matrices $\{U(q), q \in Q\}$ are unitary.

We now show that the group $\mathcal{Q} = \{U(q), q \in Q\}$ is an abelian group. First, we have that

$U(0) = UB(0)U^* = UU^* = I$ so that $I \in \mathcal{Q}$. Next, $U^{-1}(q) = UB^{-1}(q)U^* = UB(-q)U^*$ so that $U^{-1}(q) \in \mathcal{Q}$ since $-q \in \mathcal{Q}$. Finally, $U(q)U(h) = UB(q)B(h)U^* = UB(h)B(q)U^* = U(h)U(q)$ since diagonal matrices commute, and $U(q)U(h) \in \mathcal{Q}$ since $B(h)B(q) = B(h+q) = B(q')$ for some $q' \in \mathcal{Q}$.

We therefore conclude that $\phi(q) = U(q)\phi$ where the matrices $U(q)$ are unitary and form an abelian group, so that the vectors $\phi(q)$ are geometrically uniform. $\square$

As a consequence of Theorem 1 we have the following corollary.

Corollary 1. *A set of vectors $\{\phi_i \in \mathcal{H}, 1 \leq i \leq n\}$ is geometrically uniform if and only if the matrix Φ of columns ϕ_i has an SVD of the form $\Phi = U\Sigma\mathcal{F}^*$, where U is an arbitrary unitary matrix, Σ is an arbitrary diagonal matrix with diagonal elements $\{\sigma_i, 1 \leq i \leq m\}$, and $\mathcal{F}$ is a FT matrix over a direct product of cyclic groups. In addition, the vectors $\{\phi_i\}$ form a geometrically uniform frame for $\mathcal{H}$ if they span $\mathcal{H}$ or, equivalently, if $\sigma_i \neq 0$ for $1 \leq i \leq m$.*

4 Dual and Canonical Tight Frames Associated With GU Frames

In Section 4.1, we show that the dual frame vectors and the canonical tight frame vectors associated with a GU frame are also GU. This property can then be used to compute the dual and canonical tight frames very efficiently. Further properties of the canonical tight frame vectors are discussed in Section 4.2.

4.1 Constructing the dual and canonical tight frames

Let $\mathcal{S} = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}\}$ be a GU frame generated by a finite abelian group $\mathcal{Q}$ of unitary matrices, where ϕ is an arbitrary generating vector. Then the frame operator S defined by (2) commutes with each of the unitary matrices U_i in the generating group $\mathcal{Q}$. Indeed, expressing the

frame operator as

$$S = \sum_{i=1}^n U_i \phi \phi^* U_i^*, \quad (23)$$

we have that for all j ,

$$\begin{aligned} S U_j &= \sum_{i=1}^n U_i \phi_i \phi_i^* U_i^* U_j \\ &= U_j \sum_{i=1}^n U_j^* U_i \phi \phi^* U_i^* U_j \\ &= U_j \sum_{i=1}^n U_i \phi \phi^* U_i \\ &= U_j S, \end{aligned} \quad (24)$$

since $\{U_j^* U_i, 1 \leq i \leq n\}$ is just a permutation of $\mathcal{Q}$.

If S commutes with U_j , then from (5) and (8) we have that S^{-1} and $S^{-1/2}$ also commute with U_j for all j . Thus,

$$\bar{\phi}_i = S^{-1} \phi_i = S^{-1} U_i \phi = U_i S^{-1} \phi = U_i \bar{\phi}, \quad (25)$$

where $\bar{\phi} = S^{-1} \phi$, which shows that the dual frame vectors $\{\bar{\phi}_i = S^{-1} \phi_i\}$ are GU with generating group equal to $\mathcal{Q}$.

Similarly,

$$\mu_i = S^{-1/2} \phi_i = S^{-1/2} U_i \phi = U_i S^{-1/2} \phi = U_i \mu, \quad (26)$$

where $\mu = S^{-1/2} \phi$, which shows that the canonical tight frame vectors $\{\mu_i = S^{-1/2} \phi_i\}$ are also GU with generating group $\mathcal{Q}$.

Therefore, to compute the dual frame vectors or the canonical tight frame vectors all we need is to compute the generating vectors $\bar{\phi}$ and μ , respectively. The remaining frame vectors are then obtained by applying the group $\mathcal{Q}$ to the corresponding generating vectors.

We now show that the generating vectors can be computed very efficiently using the FT. From

Corollary 1 we have that Φ has an SVD of the form

$$\Phi = U\Sigma\mathcal{F}^* = \sum_{h \in Q} \sigma(h)u(h)\mathcal{F}^*(h). \quad (27)$$

Here Σ is a diagonal matrix with diagonal elements $\{\sigma(h) = n^{1/4}\sqrt{\hat{s}(h)}, h \in Q\}$ where $\{\hat{s}(h), h \in Q\}$ is the FT of $\{s(q), q \in Q\}$, U is the matrix of columns $u(h)$, where

$$u(h) = \begin{cases} \Phi\mathcal{F}(h)/\sigma(h) = \hat{\phi}(h)/\sigma(h), & \text{if } \sigma(h) \neq 0; \\ 0, & \text{otherwise} \end{cases} \quad (28)$$

with

$$\hat{\phi}(h) = \frac{1}{\sqrt{n}} \sum_{q \in Q} \langle h, q \rangle \phi(q) \quad (29)$$

denoting the h th element of the FT of Φ regarded as a row vector of column vectors, $\Phi = \{\phi(q), q \in G\}$, and $\mathcal{F}^* = \{\frac{1}{\sqrt{n}}\langle h, q \rangle^*, h, q \in Q\}$ has rows $\mathcal{F}^*(h) = \{\frac{1}{\sqrt{n}}\langle h, q \rangle^*, q \in Q\}$.

It then follows that

$$\bar{\phi} = S^{-1}\phi = \frac{1}{\sqrt{n}} \sum_{h \in \mathcal{I}} \frac{1}{\sigma(h)} u(h), \quad (30)$$

where $h \in \mathcal{I}$ if $\sigma(h) \neq 0$. Similarly,

$$\mu = S^{-1/2}\phi = \frac{1}{\sqrt{n}} \sum_{h \in \mathcal{I}} u(h). \quad (31)$$

We summarize our results in the following theorem:

Theorem 2 (GU frames). *Let $\mathcal{S} = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}\}$ be a GU frame generated by a finite abelian group $\mathcal{Q}$ of unitary matrices, where ϕ is an arbitrary generating vector, and let Φ be the matrix of columns ϕ_i . Let Q be an additive abelian group isomorphic to $\mathcal{Q}$, let $\{\phi(q), q \in Q\}$ be the elements of $\mathcal{S}$ under this isomorphism, and let $\mathcal{F}$ be the Fourier transform matrix over G . Then*

1. *the dual frame vectors $\{\bar{\phi}_i, 1 \leq i \leq n\}$ are geometrically uniform with generating group $\mathcal{Q}$ and generating vector $\bar{\phi} = (1/\sqrt{n}) \sum_{h \in \mathcal{I}} (1/\sigma(h))u(h)$, where*
 - (a) *$\{\sigma(h) = n^{1/4} \sqrt{\hat{s}(h)}, h \in Q\}$ are the singular values of Φ ,*
 - (b) *$\{\hat{s}(h), h \in Q\}$ is the Fourier transform of the inner-product sequence $\{\langle \phi(0), \phi(q) \rangle, q \in Q\}$,*
 - (c) *$\mathcal{I}$ is the set of indices $h \in Q$ for which $\sigma(h) \neq 0$,*
 - (d) *$u(h) = \hat{\phi}(h)/\sigma(h)$ for $h \in \mathcal{I}$,*
 - (e) *$\{\hat{\phi}(h), h \in Q\}$ is the Fourier transform of $\{\phi(q), q \in Q\}$,*
2. *the canonical tight frame vectors $\{\mu_i, 1 \leq i \leq n\}$ are GU with generating group $\mathcal{Q}$ and generating vector $\mu = (1/\sqrt{n}) \sum_{h \in \mathcal{I}} u(h)$;*
3. *the frame bounds of the frame $\{\phi_i, 1 \leq i \leq n\}$ are given by $A = \sqrt{n} \min_{h \in \mathcal{I}} \hat{s}(h)$ and $B = \sqrt{n} \max_{h \in \mathcal{I}} \hat{s}(h)$.*

An important special case of Theorem 2 is the case in which the generating group $\mathcal{Q}$ is *cyclic* so that $U_i = V^{i-1}, 1 \leq i \leq n$, where V is a unitary matrix with $V^n = I$. A cyclic group generates a cyclic vector set $\mathcal{S} = \{\phi_i = V^{i-1}\phi, 1 \leq i \leq n\}$, where ϕ is arbitrary. If $\mathcal{Q}$ is cyclic, then G is a circulant matrix⁵, and Q is the cyclic group $\mathbb{Z}_n$. The FT kernel is then $\langle h, g \rangle = e^{-2\pi i h g / n}$ for

⁵A circulant matrix is a matrix where every row (or column) is obtained by a right circular shift (by one position) of the previous row (or column). An example is:
$$\begin{bmatrix} a_0 & a_2 & a_1 \\ a_1 & a_0 & a_2 \\ a_2 & a_1 & a_0 \end{bmatrix}.$$

$h, g \in \mathbb{Z}_n$, and the FT matrix $\mathcal{F}$ reduces to the $n \times n$ DFT matrix. The singular values of Φ are then $n^{1/4}$ times the square roots of the DFT values of the inner products $\{\langle \phi_1, \phi_j \rangle, 1 \leq j \leq n\}$.

4.2 Properties of the canonical tight frame

The canonical tight frame vectors μ_i corresponding to the frame vectors ϕ_i have the property that they are the closest normalized tight frame vectors to the vectors ϕ_i , in a least-squares sense [28, 26, 29]. Thus, the vectors μ_i are the normalized tight frame vectors that minimize the least-squares error

$$\sum_{i=1}^n \langle \phi_i - \mu_i, \phi_i - \mu_i \rangle. \quad (32)$$

We now show that when the original frame vectors ϕ_i are GU with generating group $\mathcal{Q}$, the canonical tight frame vectors have the additional property that from all normalized tight frame vectors they maximize

$$R_{\phi\mu} = \sum_{i=1}^n |\langle \phi_i, \mu_i \rangle|^2. \quad (33)$$

Maximizing $R_{\phi\mu}$ may be of interest in various applications. For example, in a matched-filter detection problem considered in [12], $R_{\phi\mu}$ represents the total output signal-to-noise ratio. As another example, in a multiuser detection problem considered in [13], maximizing $R_{\phi\mu}$ has the effect of minimizing the multiple-access interference at the input to the proposed detector.

To obtain a more convenient expression for $R_{\phi\mu}$, let Φ and M denote the matrices of columns ϕ_i and μ_i , respectively. Since the vectors μ_i form a normalized tight frame for $\mathcal{H}$, M satisfies

$$MM^* = I_m. \quad (34)$$

From Corollary 1, Φ has an SVD of the form $\Phi = U\Sigma\mathcal{F}^*$, where U is unitary, $\mathcal{F}$ is the FT matrix over the additive group Q to which $\mathcal{Q}$ is isomorphic, and Σ is an $m \times n$ diagonal matrix with diagonal elements $\sigma_i > 0$. From (34) it follows that M can be written as $M = U\tilde{I}Z^*$ where Z is an

arbitrary unitary matrix and $\tilde{I}$ is an $m \times n$ diagonal matrix with diagonal elements all equal to 1.

Let f_i and z_i denote the columns of $\mathcal{F}^*$ and Z^* , respectively. Then we can express $R_{\phi\mu}$ as

$$R_{\phi\mu} = \sum_{i=1}^n |\langle \phi_i, \mu_i \rangle|^2 = \sum_{i=1}^n |\langle U^* \phi_i, U^* \mu_i \rangle|^2 = \sum_{i=1}^n |\langle f_i, \bar{\Sigma} z_i \rangle|^2, \quad (35)$$

where $\bar{\Sigma}$ is an $n \times n$ diagonal matrix with the first m diagonal elements equal to σ_i , and the remaining diagonal elements are all equal to 0.

Our problem then reduces to finding a set of orthonormal vectors z_i that maximize $\sum_i |\langle f_i, \bar{\Sigma} z_i \rangle|^2$, where the vectors f_i are also orthonormal. Using the Cauchy-Schwarz inequality we have that

$$R_{\phi\mu} = \sum_{i=1}^n |\langle \bar{\Sigma}^{1/2} f_i, \bar{\Sigma}^{1/2} z_i \rangle|^2 \leq \sum_{i=1}^n \langle f_i, \bar{\Sigma} f_i \rangle \langle z_i, \bar{\Sigma} z_i \rangle, \quad (36)$$

with equality if and only if $\bar{\Sigma}^{1/2} f_i = c_i \bar{\Sigma}^{1/2} z_i$ for some c_i . In particular, we have equality for $z_i = f_i$. Since the components of the vectors f_i all have equal magnitude $1/\sqrt{n}$, $\langle f_i, \bar{\Sigma} f_i \rangle = 1/n \sum_{k=1}^m \sigma_k \triangleq \alpha$ for all i , and (36) reduces to

$$R_{\phi\mu} \leq \alpha \sum_{i=1}^n \langle z_i, \bar{\Sigma} z_i \rangle = \alpha \text{Tr}(Z^* \bar{\Sigma} Z) = \alpha \text{Tr}(\bar{\Sigma}) = n\alpha^2, \quad (37)$$

with equality if $f_i = z_i$.

The normalized tight frame vectors that maximize $R_{\phi\mu}$ are then the columns of $\widehat{M} = U\tilde{I}\mathcal{F}^* = S^{-1/2}\Phi$ where $S = \Phi\Phi^*$, and are equal to the canonical tight frame vectors.

5 Example Of A GU Frame

We now consider an example demonstrating the ideas of the previous section.

Consider the frame vectors $\phi_1 = 1/2[\sqrt{3} \ -1]^*$, $\phi_2 = 1/2[\sqrt{3} \ 1]^*$, $\phi_3 = 1/2[-\sqrt{3} \ 1]^*$, $\phi_4 = 1/2[-\sqrt{3} \ -1]^*$, depicted in Fig. 1.

The corresponding Gram matrix is given by

$$G = \begin{bmatrix} 1 & 0.5 & -1 & -0.5 \\ 0.5 & 1 & -0.5 & -1 \\ -1 & -0.5 & 1 & 0.5 \\ -0.5 & -1 & 0.5 & 1 \end{bmatrix}, \quad (38)$$

which is a permuted matrix with $G = G^T$. From Proposition 2 it follows that the vectors ϕ_i are GU. Since the vectors ϕ_i also span $\mathcal{R}^2$, these vectors form a GU frame for $\mathcal{R}^2$.




Figure 1: Example of a GU frame.

The vectors ϕ_i can be expressed as $\{\phi_i = U_i\phi, 1 \leq i \leq 4\}$, where $\phi = \phi_1$ and the matrices

$\{U_i, 1 \leq i \leq 4\}$ are unitary, form an abelian group $\mathcal{Q}$, and are given by

$$U_1 = I_2, \quad U_2 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad U_3 = \begin{bmatrix} -1 & 0 \\ 0 & -1 \end{bmatrix}, \quad U_4 = \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}. \quad (39)$$

The multiplication table of the group $\mathcal{Q}$ is

	U_1	U_2	U_3	U_4
U_1	U_1	U_2	U_3	U_4
U_2	U_2	U_1	U_4	U_3
U_3	U_3	U_4	U_1	U_2
U_4	U_4	U_3	U_2	U_1

(40)

If we define the correspondence

$$U_1 \leftrightarrow (0,0), \quad U_2 \leftrightarrow (0,1), \quad U_3 \leftrightarrow (1,0), \quad U_4 \leftrightarrow (1,1), \quad (41)$$

then this table becomes the addition table of $Q = \mathbb{Z}_2 \times \mathbb{Z}_2$:

	$(0,0)$	$(0,1)$	$(1,0)$	$(1,1)$
$(0,0)$	$(0,0)$	$(0,1)$	$(1,0)$	$(1,1)$
$(0,1)$	$(0,1)$	$(0,0)$	$(1,1)$	$(1,0)$
$(1,0)$	$(1,0)$	$(1,1)$	$(0,0)$	$(0,1)$
$(1,1)$	$(1,1)$	$(1,0)$	$(0,1)$	$(0,0)$

(42)

Only the way in which the elements are labeled distinguishes the table of (42) from the table of

(40); thus $\mathcal{Q}$ is isomorphic to Q . Over $Q = \mathbb{Z}_2 \times \mathbb{Z}_2$, the FT matrix $\mathcal{F}$ is the Hadamard matrix

$$\mathcal{F} = \frac{1}{2} \begin{bmatrix} 1 & 1 & 1 & 1 \\ 1 & -1 & 1 & -1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \end{bmatrix}. \quad (43)$$

From Theorem 2 the dual frame vectors and the canonical tight frame vectors are also GU with generators $\bar{\phi}$ and μ respectively, whose equations are given in the theorem. Thus, to compute the dual and canonical tight frame vectors we compute these generators and then apply the group $\mathcal{Q}$.

We first determine the FT of the first row of G denoted by s :

$$\hat{s} = \mathcal{F}s = \frac{1}{2}[0 \ 0 \ 3 \ 1]^*. \quad (44)$$

Using Theorem 2, it follows from (44) that the frame bounds are given by $A = 1$ and $B = 3$. Next, we compute the vectors $\hat{\phi}(h)$ which are the columns of

$$\Phi\mathcal{F} = \begin{bmatrix} 0 & 0 & 1.7 & 0 \\ 0 & 0 & 0 & -1 \end{bmatrix}. \quad (45)$$

Using the equations of the theorem, we then have that $\bar{\phi} = [0.3 \ -0.5]^*$ and $\mu = [0.5 \ -0.5]^*$. By applying the group $\mathcal{Q}$ to these generators we obtain that the dual frame vectors are the columns of

$$\bar{\Phi} = \begin{bmatrix} 0.3 & 0.3 & -0.3 & -0.3 \\ -0.5 & 0.5 & 0.5 & -0.5 \end{bmatrix}, \quad (46)$$

and the canonical tight frame vectors are the columns of

$$M = \begin{bmatrix} 0.5 & 0.5 & -0.5 & -0.5 \\ -0.5 & 0.5 & 0.5 & -0.5 \end{bmatrix}. \quad (47)$$

Comparing (46) and (47) with the original frame vectors ϕ_i , it is evident that the dual and canonical tight frame vectors have the same symmetries as the original frame vectors, as illustrated in Fig. 2.




Figure 2: Symmetry property of the frame vectors ϕ_i , the dual frame vectors $\bar{\phi}_i$, and the canonical tight frame vectors μ_i . $\bar{\phi}_i$ are the columns of $\bar{\Phi}$ given by (46), and μ_i are the columns of M given by (47). The frame vectors, dual frame vectors, and the canonical tight frame vectors all have the same symmetry properties.

6 Compound GU Frames

In Section 4 we showed that the dual and canonical tight frame vectors associated with a GU frame are themselves GU and can therefore be computed using a single generator. In this section, we consider a class of frames which consist of subsets that are GU, and are therefore referred to as *compound geometrically uniform (CGU) frames*. As we show, the dual and canonical tight frame vectors associated with a CGU frame share the same symmetries as the original frame and can be computed using a *set* of generators.

A set of frame vectors $\{\phi_{ik}, 1 \leq i \leq l, 1 \leq k \leq r\}$ is CGU if $\phi_{ik} = U_i \phi_k$ for some generating vectors $\{\phi_k, 1 \leq k \leq r\}$, and the matrices $\{U_i, 1 \leq i \leq l\}$ are unitary and form an abelian group $\mathcal{Q}$.

A CGU frame is in general not GU. However, for every k , the vectors $\{\phi_{ik}, 1 \leq i \leq l\}$ are a GU vector set with generating group $\mathcal{Q}$.

A special case of CGU frames are filter bank frames studied in [7, 8, 9], in which $\mathcal{Q}$ is the group of translations by integer multiples of the subsampling factor, and the generating vectors are the filter bank synthesis filters. We may therefore view CGU frames as an extension of filter bank frames to the more general group case.

As we show in the following proposition, the frame bounds of a CGU frame can be bounded by the sum of the norms of the generating vectors.

Proposition 3. *Let $\mathcal{S} = \{\phi_{ik} = U_i \phi_k, 1 \leq i \leq l, 1 \leq k \leq r\}$ be a compound geometrically uniform frame with frame bounds A and B , where $\{\phi_k, 1 \leq k \leq r\}$ is an arbitrary set of generating vectors. Then $A \leq \frac{l}{m} \sum_{k=1}^r \|\phi_k\|^2 \leq B$. If in addition the frame is tight, then $A = \frac{l}{m} \sum_{k=1}^r \|\phi_k\|^2$.*

Proof. We can express the frame operator corresponding to the frame vectors ϕ_i as

$$S = \sum_{i=1}^l \sum_{k=1}^r U_i \phi_k \phi_k^* U_i^*. \quad (48)$$

Then,

$$\mathrm{Tr}(S) = \sum_{i=1}^l \sum_{k=1}^r \mathrm{Tr}(U_i \phi_k \phi_k^* U_i^*) = l \sum_{k=1}^r \|\phi_k\|^2, \quad (49)$$

so that

$$\sum_{i=1}^m \lambda_i(S) = \mathrm{Tr}(S) = l \sum_{k=1}^r \|\phi_k\|^2. \quad (50)$$

Therefore,

$$A = \min_i \lambda_i(S) \leq \frac{1}{m} \sum_{i=1}^m \lambda_i(S) = \frac{l}{m} \sum_{k=1}^r \|\phi_k\|^2, \quad (51)$$

and

$$B = \max_i \lambda_i(S) \geq \frac{1}{m} \sum_{i=1}^m \lambda_i(S) = \frac{l}{m} \sum_{k=1}^r \|\phi_k\|^2. \quad (52)$$

□

6.1 Dual and canonical tight frames associated with CGU frames

We now show that the dual and canonical tight frames associated with a CGU frame are also CGU.

Expressing the frame operator as

$$S = \sum_{i=1}^l \sum_{k=1}^r \phi_{ik} \phi_{ik}^* = \sum_{i=1}^l U_i \left(\sum_{k=1}^r \phi_k \phi_k^* \right) U_i^*, \quad (53)$$

for all j we have that,

$$\begin{aligned} S U_j &= \sum_{i=1}^l U_i \left(\sum_{k=1}^r \phi_k \phi_k^* \right) U_i^* U_j \\ &= U_j \sum_{i=1}^l U_j^* U_i \left(\sum_{k=1}^r \phi_k \phi_k^* \right) U_i^* U_j \\ &= U_j \sum_{i=1}^l U_i \left(\sum_{k=1}^r \phi_k \phi_k^* \right) U_i^* \\ &= U_j S, \end{aligned} \quad (54)$$

since $\{U_j^* U_i, 1 \leq i \leq l\}$ is just a permutation of $\mathcal{Q}$. Thus S commutes with U_j , so that S^{-1} and $S^{-1/2}$ also commute with U_j for all j . Then, the dual frame vectors $\bar{\phi}_{ik}$ of the vectors ϕ_{ik} are given by

$$\bar{\phi}_{ik} = S^{-1} \phi_{ik} = S^{-1} U_i \phi_k = U_i S^{-1} \phi_k = U_i \bar{\phi}_k, \quad (55)$$

where $\bar{\phi}_k = S^{-1} \phi_k$, which shows that the dual frame vectors $\{\bar{\phi}_{ik} = S^{-1} \phi_{ik}\}$ are CGU with generating group equal to $\mathcal{Q}$.

Similarly,

$$\mu_{ik} = S^{-1/2} \phi_{ik} = S^{-1/2} U_i \phi_k = U_i S^{-1/2} \phi_k = U_i \mu_k, \quad (56)$$

where $\mu_k = S^{-1/2} \phi_k$, which shows that the canonical tight frame vectors $\{\mu_{ik} = S^{-1/2} \phi_{ik}\}$ are also CGU with generating group $\mathcal{Q}$.

Therefore, to compute the dual frame vectors or the canonical tight frame vectors all we need is to compute the generating vectors $\{\bar{\phi}_k, 1 \leq k \leq r\}$ and $\{\mu_k, 1 \leq k \leq r\}$, respectively. The remaining frame vectors are then obtained by applying the group $\mathcal{Q}$ to the corresponding set of generating vectors.

6.2 CGU frames with GU generators

A special class of CGU frames is *CGU frames with GU generators* in which the generating vectors $\{\phi_k, 1 \leq k \leq r\}$ are themselves GU. Specifically, $\{\phi_k = V_k \phi\}$ for some generator ϕ , where the matrices $\{V_k, 1 \leq k \leq r\}$ are unitary, and form an abelian group $\mathcal{G}$.

Now suppose that U_p and V_t commute up to a phase factor for all t and p so that $U_p V_t = V_t U_p e^{j\theta(p,t)}$ where $\theta(p,t)$ is an arbitrary phase function that may depend on the indices p and t . In this case we say that $\mathcal{Q}$ and $\mathcal{G}$ commute up to a phase factor. Then for all p, t ,

$$S U_p V_t = \sum_{i=1}^l U_i \left(\sum_{k=1}^r V_k \phi \phi^* V_k^* \right) U_i^* U_p V_t$$

$$\begin{aligned}
&= U_p V_t \sum_{i=1}^l V_t^* U_p^* U_i \left(\sum_{k=1}^r V_k \phi \phi^* V_k^* \right) U_i^* U_p V_t \\
&= U_p V_t \sum_{i=1}^l V_t^* U_i^* \left(\sum_{k=1}^r V_k \phi \phi^* V_k^* \right) U_i V_t \\
&= U_p V_t \sum_{i=1}^l U_i^* \left(\sum_{k=1}^r V_t^* V_k \phi \phi^* V_k^* V_t \right) U_i \\
&= U_p V_t \sum_{i=1}^l U_i \left(\sum_{k=1}^r V_k \phi \phi^* V_k^* \right) U_i^* \\
&= U_p V_t S.
\end{aligned} \tag{57}$$

The dual frame vectors $\bar{\phi}_{ik}$ of the vectors ϕ_{ik} are then given by

$$\bar{\phi}_{ik} = S^{-1} \phi_{ik} = S^{-1} U_i V_k \phi = U_i V_k S^{-1} \phi = U_i V_k \bar{\phi}, \tag{58}$$

where $\bar{\phi} = S^{-1} \phi$. Similarly,

$$\mu_{ik} = S^{-1/2} \phi_{ik} = S^{-1/2} U_i V_k \phi = U_i V_k S^{-1/2} \phi = U_i V_k \mu \tag{59}$$

where $\mu = S^{-1/2} \phi$. Thus even though the frame is not in general GU, the dual and canonical tight frame vectors can be computed using a single generating vector.

As we now show, in the special case in which $\theta = 0$ so that $U_i V_k = V_k U_i$ for all i, k , the resulting frame is GU. To this end we need to show that the unitary matrices $\mathcal{Q}' = \{Q_{ik} = U_i V_k\}$ form an abelian group. First, $Q_{ik} Q_{jt} = U_i V_k U_j V_t = U_i U_j V_k V_t$. Since $U_i U_j \in \mathcal{Q}$ and $V_k V_t \in \mathcal{G}$, $Q_{ik} Q_{jt} \in \mathcal{Q}'$. Next, $Q_{ik}^{-1} = V_k^* U_i^* = U_i^* V_k^* \in \mathcal{Q}'$ since $U_i^* \in \mathcal{Q}$ and $V_k^* \in \mathcal{G}$. Also, $I \in \mathcal{Q}'$ since $I \in \mathcal{Q}$ and $I \in \mathcal{G}$. Finally, $Q_{ik} Q_{jt} = U_i V_k U_j V_t = U_i U_j V_t V_k = U_j V_t U_i V_k = Q_{jt} Q_{ik}$.

A special case of CGU frames with GU generators for which $\mathcal{Q}$ and $\mathcal{G}$ commute up to a phase factor are Weyl-Heisenberg (WH) frames [17, 3, 16]. If the WH frame is critically sampled, then $\theta(p, t) = 0$ and the WH frame reduces to a GU frame. In the more general oversampled case,

$\theta(p, t) \neq 0$. CGU frames with GU generators can hence be viewed as a generalization of WH frames to the group case.

To summarize, we have the following theorem:

Theorem 3 (Compound GU frames). *Let $\mathcal{S} = \{\phi_{ik} = U_i\phi_k, 1 \leq i \leq l, 1 \leq k \leq r\}$ be a compound geometrically uniform frame with generating vectors $\{\phi_k, 1 \leq k \leq r\}$ and generating group $\mathcal{Q}$, and let S be the frame operator corresponding to the frame vectors $\{\phi_{ik}\}$. Then*

1. *the dual frame vectors $\{\bar{\phi}_{ik}, 1 \leq i \leq l, 1 \leq k \leq r\}$ are compound geometrically uniform with generating group $\mathcal{Q}$ and generating vectors $\{\bar{\phi}_k = S^{-1}\phi_k, 1 \leq k \leq r\}$;*
2. *the canonical tight frame vectors $\{\mu_{ik}, 1 \leq i \leq l, 1 \leq k \leq r\}$ are compound geometrically uniform with generating group $\mathcal{Q}$ and generating vectors $\{\mu_k = S^{-1/2}\phi_k, 1 \leq k \leq r\}$.*

If in addition the generating vectors $\{\phi_k = V_k\phi, 1 \leq k \leq r\}$ are geometrically uniform with $U_iV_k = V_kU_ie^{j\theta(i,k)}$ for all i, k , then

1. *$\bar{\phi}_{ik} = U_iV_k\bar{\phi}$ where $\bar{\phi} = S^{-1}\phi$;*
2. *$\mu_{ik} = U_iV_k\mu$ where $\mu = S^{-1/2}\phi$;*
3. *if in addition $\theta(i, k) = 0$ for all i, k , then the vectors $\{\phi_{ik}, 1 \leq i \leq l, 1 \leq k \leq r\}$ form a geometrically uniform frame.*

7 Pruning GU Frames

In applications it is often desirable to know how a frame behaves when one or more frame elements are removed. In particular, it is important to know or to be able to estimate the frame bounds of the reduced frame. In general, if no structural constraints are imposed on a frame this behavior will depend critically on the particular frame elements removed. For example, removal of a particular frame element may destroy the frame property so that the remaining vectors do not constitute a frame anymore, whereas if a different element is removed the remaining vectors may still constitute a frame.

One of the prime applications of frames is signal analysis and synthesis, where a signal is expanded by computing the inner products of the signal with the frame elements. The resulting coefficients are subsequently stored, transmitted, quantized or manipulated in some way. In particular, a coefficient may be lost (*e.g.*, due to a transmission error) which results in a reconstructed signal that is equivalent to an expansion using a pruned frame obtained by removing the corresponding frame vector.

Recently, there has been increased interest in using frames for multiple-description source coding where a signal is expanded into a redundant set of functions and the resulting coefficients are transmitted over a lossy packet network where one or more of the coefficients can be lost because a packet is dropped [14]. The goal of multiple description source coding is to ensure a gradually behaving reconstruction quality as a function of the number of dropped packets. When using frames in this context, the reconstruction quality is often governed by the frame bound ratio of the pruned frame. If the packets are dropped with equal probability, then it is desirable that the frame bound ratio should deteriorate uniformly irrespectively of the particular frame element that is removed. In the following, we show that GU frames have this property. We will furthermore demonstrate that if the original frame is a tight GU frame, then the frame bound ratio of the pruned frame obtained by removing one frame element can be computed exactly. We also consider the case where sets of

frame elements are removed.

Theorem 4 (Pruned GU frames). *Let $\mathcal{S} = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}\}$ be a geometrically uniform frame generated by a finite abelian group $\mathcal{Q}$ of unitary matrices, where ϕ is an arbitrary generating vector. Let Φ be the matrix of columns ϕ_i , and let $S = \Phi\Phi^*$ be the corresponding frame operator. Let $\mathcal{S}(j) = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}, i \neq j\}$ be the pruned set obtained by removing the element ϕ_j . Then the eigenvalues of the frame operator corresponding to the pruned set do not depend on the particular element ϕ_j removed.*

Proof. The frame operator corresponding to the pruned frame is given by

$$S(j) = \sum_{i=1}^n U_i \phi \phi^* U_i^* - U_j \phi \phi^* U_j^*. \quad (60)$$

Since U_j is unitary, the eigenvalues of $S(j)$ are equal to the eigenvalues of $U_j^* S(j) U_j$. But,

$$U_j^* S(j) U_j = U_j^* \sum_{i=1}^n U_i \phi \phi^* U_i^* U_j - \phi \phi^* = \sum_{i=1}^n U_i \phi \phi^* U_i^* - \phi \phi^* = S - \phi \phi^*. \quad (61)$$

Since $U_j^* S(j) U_j$ is independent of j , the eigenvalues of $S(j)$ do not depend on j . $\square$

In general it is difficult to provide estimates on the frame bounds of the pruned frame. However, in the special case where the original GU frame is a tight frame, these bounds can be determined exactly.

Corollary 2 (Pruned Tight GU frames). *Let $\mathcal{S} = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}\}$ be a geometrically uniform tight frame generated by a finite abelian group $\mathcal{Q}$ of unitary matrices, where ϕ is a unit norm generating vector. Let Φ be the matrix of columns ϕ_i , and let $S = \Phi\Phi^*$ be the corresponding frame operator. Let $\mathcal{S}(j) = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}, i \neq j\}$ be the pruned set obtained by removing the element ϕ_j . Then the eigenvalues of the frame operator corresponding to the pruned set are given by $\lambda_1 = \frac{n}{m} - 1$ and $\lambda_i = \frac{n}{m}$, $2 \leq i \leq n$, independent of ϕ_j .*

Proof. Since $\mathcal{S}$ is a tight frame with $\|\phi\| = 1$, from Proposition 1 the frame bound $A = n/m$ and $S = (n/m)I_m$. Then,

$$S(j) = \frac{n}{m}I_m - U_j\phi\phi^*U_j^*, \quad (62)$$

and

$$U_j^*S(j)U_j = \frac{n}{m}I_m - \phi\phi^*. \quad (63)$$

Since $\|\phi\| = 1$, $\phi\phi^*$ has one eigenvalue equal to 1, and the remaining eigenvalues equal to 0. The eigenvalues of $S(j)$ are therefore given by $\lambda_1 = \frac{n}{m} - 1$ and $\lambda_i = \frac{n}{m}$, $2 \leq i \leq n$. $\square$

An immediate consequence of Corollary 2 is that the frame bound ratio of the pruned frame is given by $B/A = 1/(1 - m/n)$, which is close to 1 for large redundancy $r = n/m$.

We next consider the case where multiple frame elements are removed.

Corollary 3. *Let $\mathcal{S} = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}\}$ be a geometrically uniform frame generated by a finite abelian group $\mathcal{Q}$ of unitary matrices, let Φ be the matrix of columns ϕ_i , and let $S = \Phi\Phi^*$ be the corresponding frame operator. Let $\mathcal{J}$ be a set of indices, and let $\mathcal{J}(k)$ denote the set of indices i such that $U_i = U_kU_j$ for fixed k and $j \in \mathcal{J}$. Let $\mathcal{S}(k) = \{\phi_i = U_i\phi, U_i \in \mathcal{Q}, i \notin \mathcal{J}(k)\}$ be a pruned set obtained by removing the elements ϕ_i with $i \in \mathcal{J}(k)$. Then the eigenvalues of the frame operator corresponding to the pruned set are independent of k .*

Proof. The frame operator corresponding to the pruned frame is given by

$$S(k) = \sum_{i=1}^n U_i\phi\phi^*U_i^* - U_k \left[\sum_{j \in \mathcal{J}} U_j\phi\phi^*U_j^* \right] U_k^*. \quad (64)$$

Then,

$$U_k^*S(k)U_k = U_k^* \sum_{i=1}^n U_i\phi\phi^*U_i^*U_k - \sum_{j \in \mathcal{J}} U_j\phi\phi^*U_j^*$$

$$= \sum_{i=1}^n U_i \phi \phi^* U_i^* - \sum_{j \in \mathcal{J}} U_j \phi \phi^* U_j^*$$

is independent of k , and consequently the eigenvalues of $S(k)$ do not depend on k . $\square$

To conclude this section, GU frames have strong symmetry properties in the sense that removing any one of the elements leads to a vector set with bounds independent of the particular element removed. Moreover, if the original frame is tight, then we can compute the bounds of the pruned frame exactly.

8 Constructing GU Frames

Suppose we are given a set of vectors $\{\varphi_i, 1 \leq i \leq n\}$, that form a frame for an m -dimensional space $\mathcal{H}$. We would like to construct a geometrically uniform frame $\{\phi_i, 1 \leq i \leq n\}$ from the vectors $\{\varphi_i\}$.

From Theorem 1 it follows that the vectors $\{\phi_i\}$ form a GU frame if and only if the Gram matrix G has rank m , and is diagonalized by a FT matrix $\mathcal{F}$ over a finite product of cyclic groups. There are many ways to construct a frame from a given set of frame vectors φ_i that satisfy these properties. For example, let F be the matrix of columns φ_i , and let F have an SVD $F = Q\Lambda V^*$, where Λ is a diagonal matrix with diagonal elements λ_i . Then the columns of $\Phi = FV\Sigma\mathcal{F}^* = Q\Lambda\Sigma\mathcal{F}^*$ form a GU frame, where $\mathcal{F}$ is any FT matrix over a product of cyclic groups, and Σ is an arbitrary diagonal matrix with diagonal elements $\sigma_i > 0$. The frame bounds of the resulting GU frame are given by $A = \min_i \lambda_i^2 \sigma_i^2$ and $B = \max_i \lambda_i^2 \sigma_i^2$, so that we can choose the diagonal matrix Σ to control these bounds. In particular, choosing $\Sigma = I$ we have that the columns of $\Phi = FV\mathcal{F}^* = Q\Lambda\mathcal{F}^*$ form a GU frame. This choice has the property that the frame bounds of the GU frame are equal to the frame bounds of the original frame.

We now consider the problem of constructing an *optimal* GU frame. Specifically, let $\{\varphi_i, 1 \leq$

$i \leq n\}$ denote a frame for $\mathcal{H}$, and suppose we wish to construct a GU frame $\{\phi_i\}$ from the vectors $\{\varphi_i\}$. A reasonable approach is to find a set of vectors ϕ_i that span $\mathcal{H}$, and are “closest” to the vectors φ_i in the least-squares sense. Thus we seek vectors ϕ_i that minimize the least-squares error E , defined by

$$E = \sum_{i=1}^n \langle e_i, e_i \rangle, \quad (65)$$

where e_i denotes the i th error vector

$$e_i = \varphi_i - \phi_i, \quad (66)$$

subject to the constraint that the vectors ϕ_i form a GU frame.

If the vectors ϕ_i are GU, then their Gram matrix $G = \Phi^* \Phi$ is a permuted matrix with rank m , diagonalized by a FT matrix $\mathcal{F}$. Thus, the inner products $\{\langle \phi_i, \phi_j \rangle\}$ must satisfy

$$\{\langle \phi_i, \phi_j \rangle, 1 \leq j \leq n\} = \beta^2 P_i \{a_j, 1 \leq j \leq n\}, \quad (67)$$

where $P_i \{a_j, 1 \leq j \leq n\}$ is a permutation of the numbers $\{a_j, 1 \leq j \leq n\}$, $\beta > 0$ is a scaling factor, and the numbers $\{a_j, 1 \leq j \leq n\}$ are chosen such that the matrix R whose i th row is equal to $P_i \{a_j, 1 \leq j \leq n\}$ is Hermitian, non-negative definite and diagonalized by a FT matrix $\mathcal{F}$.

In our development of the optimal GU frame vectors we assume that the permutations P_i in (67) are specified. Since these permutations determine the additive group Q over which the FT matrix $\mathcal{F}$ is defined, we assume that $\mathcal{F}$ is specified. We then consider three different constraints on the vectors ϕ_i . First we consider the case in which both the numbers $\{a_j, 1 \leq j \leq n\}$ and the scaling β in (67) are known. The GU frame minimizing the least-squares error E of (65)-(66) subject to this constraint is derived in Section 8.1, and is referred to as the scaled-constrained least-squares GU frame (SC-LSGUF). Next, we consider the case in which the numbers $\{a_j, 1 \leq j \leq n\}$ in (67) are known, and the scaling β is chosen to minimize E . The resulting GU frame is referred to as the constrained least-squares GU frame (C-LSGUF), and is derived in Section 8.2. Finally,

in Section 8.3 we consider the more general case in which both the numbers $\{a_j, 1 \leq j \leq n\}$ and the scaling β in (67) are chosen to minimize E . The resulting GU frame is referred to as the least-squares GU frame (LSGUF).

8.1 Scaled-constrained least-squares GU frame

We first consider the case in which the numbers $\{a_j, 1 \leq j \leq n\}$ and the scaling β in (67) are known. Thus, we seek the set of vectors $\{\phi_i\}$ that minimize the least-squares error E of (65)-(66) subject to the constraint

$$\Phi^* \Phi = \beta_0^2 R = \beta_0^2 \mathcal{F} A \mathcal{F}^*, \quad (68)$$

where Φ is the matrix of columns ϕ_i , β_0 is a known scaling factor, and R is the matrix whose i th row is equal to $P_i\{a_j, 1 \leq j \leq n\}$ where the numbers $\{a_j, 1 \leq j \leq n\}$ are given such that R is diagonalized by $\mathcal{F}$, and A is a diagonal matrix with diagonal elements $\{\alpha_j = n^{1/2} \hat{a}_j, 1 \leq j \leq n\}$ where $\{\hat{a}_j, 1 \leq j \leq n\}$ is the FT of the sequence $\{a_j, 1 \leq j \leq n\}$. From (68), the frame bounds of the vectors $\{\phi_i\}$ are given by $A = \beta_0^2 \min_j \alpha_j$ and $B = \beta_0^2 \max_j \alpha_j$.

This problem has been solved in the context of general least-squares inner product shaping [37]. The optimal SC-LSGUF vectors $\hat{\phi}_i$ are the columns of $\hat{\Phi}$, given by

$$\hat{\Phi} = \beta_0 U V^* \Sigma \mathcal{F}^*, \quad (69)$$

where U and V are the right-hand unitary matrix and left-hand unitary matrix respectively in the SVD of $F \mathcal{F} \Sigma^*$, F is the matrix of columns φ_i , and Σ is an $m \times n$ diagonal matrix with diagonal elements $\sqrt{\alpha_i}$ for values of i for which $\alpha_i \neq 0$.

If $F R F^* = \beta_0^2 F \mathcal{F} A \mathcal{F}^* F$ is invertible, then we may express $\hat{\Phi}$ as

$$\hat{\Phi} = \beta_0 (F \mathcal{F} A \mathcal{F}^* F)^{-1/2} F \mathcal{F} A \mathcal{F}^* = \beta_0 (F R F^*)^{-1/2} F R. \quad (70)$$

8.2 Constrained least-squares GU frame

We now consider the case in which the numbers $\{a_j, 1 \leq j \leq n\}$ are known, but the scaling β is not specified. Thus, we seek a set of vectors $\{\phi_i\}$ that minimize the least-squares error E subject to

$$\Phi^* \Phi = \beta^2 R, \quad (71)$$

where $\beta > 0$.

The least-squares error E of (65)-(66) may be expressed as

$$\begin{aligned} E &= \text{Tr}((\Phi - F)^*(\Phi - F)) \\ &= \text{Tr}(\Phi^* \Phi) + \text{Tr}(F^* F) - 2\Re\{\text{Tr}(F^* \Phi)\} \\ &= \beta^2 \text{Tr}(R) + \text{Tr}(F^* F) - 2\Re\{\text{Tr}(F^* \Phi)\}. \end{aligned} \quad (72)$$

Let $\tilde{\Phi} = (1/\beta)\Phi$. Then minimizing E is equivalent to minimizing

$$E' = \beta^2 \text{Tr}(R) - 2\beta \Re\{\text{Tr}(F^* \tilde{\Phi})\}, \quad (73)$$

subject to

$$\tilde{\Phi}^* \tilde{\Phi} = R. \quad (74)$$

To determine the optimal matrix Φ we have to minimize E' with respect to β and $\tilde{\Phi}$. Fixing β and minimizing with respect to $\tilde{\Phi}$, the optimal value of $\tilde{\Phi}$ is given by the SC-LSGUF of Section 8.1 with scaling $\beta_0 = 1$, so that

$$\hat{\tilde{\Phi}} = UV^* \Sigma \mathcal{F}^*. \quad (75)$$

If FRF^* is invertible, then

$$\widehat{\Phi} = (F\mathcal{F}A\mathcal{F}^*F^*)^{-1/2}F\mathcal{F}A\mathcal{F}^* = (FRF^*)^{-1/2}FR. \quad (76)$$

Substituting $\widehat{\Phi}$ back into (73), and minimizing with respect to β , the optimal value of β is given by

$$\widehat{\beta} = \frac{\Re\{\text{Tr}(F^*\widehat{\Phi})\}}{\text{Tr}(R)} = \frac{\text{Tr}(UV^*\Sigma\mathcal{F}^*)}{\text{Tr}(R)}, \quad (77)$$

which in the case that FRF^* is invertible reduces to

$$\widehat{\beta} = \frac{\text{Tr}((FRF^*)^{1/2})}{\text{Tr}(R)}. \quad (78)$$

The C-LSGUF vectors are then the columns of $\widehat{\Phi}$ given by

$$\widehat{\Phi} = \widehat{\beta}UV^*\Sigma\mathcal{F}^*, \quad (79)$$

where $\widehat{\beta}$ is given by (77). If FRF^* is invertible, then

$$\widehat{\Phi} = \widehat{\beta}(FRF^*)^{-1/2}FR, \quad (80)$$

where $\widehat{\beta}$ is given by (78).

We summarize our results regarding constrained optimal GU frames in the following theorem:

Theorem 5 (Constrained least-squares GU frames). *Let $\{\varphi_i\}$ be a set of n vectors in an m -dimensional Hilbert space $\mathcal{H}$ that span $\mathcal{H}$, and let F be the matrix of columns φ_i . Let $\{\hat{\phi}_i\}$ denote the optimal n GU frame vectors that minimize the least-squares error defined by (65)-(66), subject to the constraint (67), and let $\hat{\Phi}$ be the matrix of columns ϕ_i . Let R be the matrix with i th row equal to $P_i\{a_j, 1 \leq j \leq n\}$, so that $\beta^2 R$ is the Gram matrix of inner products $\{\langle \phi_i, \phi_j \rangle\}$, and let $\mathcal{F}$ be the Fourier transform matrix that diagonalizes R . Let A be the diagonal matrix with diagonal elements $\{\alpha_j = n^{1/2} \hat{a}_j, 1 \leq j \leq n\}$ where $\{\hat{a}_j, 1 \leq j \leq n\}$ is the FT of the sequence $\{a_j, 1 \leq j \leq n\}$, let Σ be an $m \times n$ diagonal matrix with diagonal elements $\sqrt{\alpha_i}$ for values of i for which $\alpha_i \neq 0$, and let U and V be the right-hand unitary matrix and left-hand unitary matrix respectively in the SVD of $F\mathcal{F}\Sigma^*$. Then,*

1. *if $\beta = \beta_0$ is given, then $\hat{\Phi} = \beta_0 UV^* \Sigma \mathcal{F}^*$. If in addition FRF^* is invertible, then $\hat{\Phi} = \beta_0 (F\mathcal{F}A\mathcal{F}^*F^*)^{-1/2} F\mathcal{F}A\mathcal{F}^* = \beta_0 (FRF^*)^{-1/2} FR$;*
2. *if $\beta > 0$ is chosen to minimize E , then $\hat{\Phi} = \hat{\beta} UV^* \Sigma \mathcal{F}^*$, where $\hat{\beta}$ is given by (77). If in addition FRF^* is invertible, then $\hat{\Phi} = \hat{\beta} (F\mathcal{F}A\mathcal{F}^*F^*)^{-1/2} F\mathcal{F}A\mathcal{F}^* = \hat{\beta} (FRF^*)^{-1/2} FR$, where $\hat{\beta}$ is given by (78).*

8.3 Least-squares GU frame

We now consider the least-squares problem in which both the scaling factor β and the numbers $\{a_j, 1 \leq j \leq n\}$ in (67) are chosen to minimize E . Thus, we seek a set of vectors $\{\phi_i\}$ that minimize the least-squares error E of (65)-(66) subject to

$$\{\langle \phi_i, \phi_j \rangle, 1 \leq j \leq n\} = P_i \{a_j, 1 \leq j \leq n\}, \quad (81)$$

where $P_i \{a_j, 1 \leq j \leq n\}$ is a known permutation of the unknown numbers $\{a_j, 1 \leq j \leq n\}$, chosen such that the matrix R whose i th row is equal to $P_i \{a_j, 1 \leq j \leq n\}$ is Hermitian, non-negative definite and diagonalized by a FT matrix $\mathcal{F}$.

This problem has also been considered in the context of general least-squares inner product shaping [37]. It can be shown that the solution involves solving a problem of the form

$$\max \sum_{i=1}^m |\langle x_i, y_i \rangle|^2, \quad (82)$$

subject to

$$\langle y_i, y_j \rangle = \delta_{ij}, \quad (83)$$

where the vectors $\{x_i\}$ are known and are a function of the given vectors $\{\varphi_i\}$.

As we now show, this problem is equivalent to a quantum detection problem, for which there is no known analytical solution in the general case.

8.3.1 Connection with quantum detection

In a quantum detection problem, a system is prepared in one of m known (pure) states that are described by vectors s_i in a Hilbert space $\mathcal{H}$, and the problem is to detect the state prepared by performing a measurement on the system. The measurement is described in terms of a set of

orthogonal measurement vectors q_i . Given a set of measurement vectors q_i , and assuming equal prior probabilities on the different states, the probability of detection is given by [38],

$$P_{qd} = \frac{1}{m} \sum_{i=1}^m |\langle q_i, s_i \rangle|^2. \quad (84)$$

Comparing (84) with (82) we see that finding a set of orthogonal measurement vectors to maximize the probability of detection is equivalent to the maximization problem of (82)-(83).

Necessary and sufficient conditions for an optimum measurement maximizing (84) have been derived [39, 40, 38]. However, except in some particular cases [38, 41, 42], obtaining a closed-form analytical expression for the optimal measurement directly from these conditions is a difficult and unsolved problem. Iterative algorithms for maximizing (84) for arbitrary vector sets are given in [43, 37].

We conclude that in general there is no known analytical expression for the GULSF. In practice, the GULSF may be obtained using the iterative algorithms of [43, 37]. A more detailed discussion on the GULSF can be found in [37].

9 Distance Properties of GU Frames

So far we have mainly been concerned with structural properties of GU frames. In this section we study the Euclidean distance properties of GU frames.

Suppose we are given a GU frame $\{\phi_i = U_i\phi, 1 \leq i \leq n\}$ generated by the group $\mathcal{Q}$ with $\|\phi\|^2 = 1$. We would like to characterize the distance profile $\alpha(i, j) = \|\phi_i - \phi_j\|^2$ for all i, j .

Since the vectors ϕ_i are geometrically uniform, $\{\alpha(i, j), 1 \leq j \leq n\}$ is just a permutation of $\{d(i) = \|\phi - \phi_i\|^2, 1 \leq i \leq n\}$. Furthermore,

$$d(i) = \langle \phi, \phi \rangle + \langle U_i\phi, U_i\phi \rangle - 2\Re(\langle U_i\phi, \phi \rangle) = 2(1 - \Re(a_i)), \quad (85)$$

where $\{a_i, 1 \leq i \leq n\}$ are the elements of the first row of the Gram matrix corresponding to the frame $\{\phi_i\}$.

In applications it may be desirable to construct a GU frame such that $d(i) > 0$ for $2 \leq i \leq n$. Since $d(i) = \|(I - U_i)\phi\|^2$, a sufficient condition is that

$$\det(I - U_i) \neq 0, \quad 2 \leq i \leq n, \quad (86)$$

which is satisfied if and only if none of the matrices U_i has an eigenvalue equal to 1. Note that if (86) is satisfied, then $d(i) > 0, 2 \leq i \leq n$ regardless of the generating vector $\phi \neq 0$. Groups with unitary representations satisfying (86) are known as *fixed-point free groups*, and have been studied extensively in the literature (see *e.g.*, [44]). Thus if $\mathcal{Q}$ is a representation of a fixed-point free group, then we have that $d(i) > 0, 2 \leq i \leq n$.

Fixed-point free groups have recently been studied in the context of unitary space-time codes [33]. In particular, it was shown in [33] that an abelian group of matrices $\{U_i\}$ satisfies (86) if and only if it is cyclic, *i.e.*, $U_i = U^i$ with $U^n = I$, and where U can be parameterized as

$$U = \text{diag}(e^{j2\pi u_1/n}, \dots, e^{j2\pi u_n/n}), \quad (87)$$

where u_k is relatively prime to n for all k . An optimization over the u_k can be performed to obtain distance profiles with certain prescribed properties (there are $\phi(n)$ positive integers less than n that are relatively prime to n , where $\phi(n)$ denotes the Euler totient function of n).

10 Conclusion

In this paper we introduced the concept of GU and CGU frames and discussed some of their key properties. These frames may be viewed as generalizations of WH frames and filter bank frames to the group case. A fundamental characteristic of these frames is that they possess strong symmetry properties that may be desirable in a variety of applications. In particular, like WH frames and wavelet frames, GU frames are generated by a single generating vector. Furthermore, the canonical and dual frame vectors associated with a GU frame are themselves GU and are therefore also generated by a single generating vector which can be computed very efficiently using a Fourier transform matrix defined over an appropriate group.

We also showed that GU frame vectors possess interesting symmetry properties when one or more frame elements are removed. This property of GU frames may be of importance in multiple description source-coding where it is often desirable that the quality of the reconstruction should not depend on the particular element lost (removed).

Although in this paper we have focused on the case in which the underlying group is a finite abelian group, many of the results can be extended to the more general case of infinite-dimensional and non-abelian groups. An interesting direction for further research is to characterize these more general cases of GU frames using possibly continuous-time Fourier transforms defined over non-abelian groups (see *e.g.*, [45]).

Acknowledgments

The authors wish to thank Prof. G. D. Forney for many fruitful discussions.

References

- [1] R. J. Duffin and A. C. Schaeffer, “A class of nonharmonic Fourier series,” *Trans. Amer. Math. Soc.*, vol. 72, pp. 314–366, 1952.
- [2] R. M. Young, *An introduction to nonharmonic Fourier series*, New York: Academic Press, 1980.
- [3] C. E. Heil and D. F. Walnut, “Continuous and discrete wavelet transforms,” *SIAM Rev.*, vol. 31, no. 4, pp. 628–666, Dec. 1989.
- [4] A. Aldroubi, “Portraits of frames,” *Proc. Amer. Math. Soc.*, vol. 123, pp. 1661–1668, 1995.
- [5] J. J. Benedetto, “Irregular sampling and frames,” in *Wavelets — A tutorial in theory and applications*, C. K. Chui, Ed., pp. 445–507. Boca Raton, FL: CRC Press, 1992.
- [6] I. Daubechies, “The wavelet transform, time-frequency localization and signal analysis,” *IEEE Trans. Inform. Theory*, vol. 36, pp. 961–1005, Sep. 1990.
- [7] H. Bölcskei, *Oversampled filter banks and predictive subband coders*, Ph.D. thesis, Vienna University of Technology, Nov. 1997.
- [8] H. Bölcskei, F. Hlawatsch, and H. G. Feichtinger, “Frame-theoretic analysis of oversampled filter banks,” *IEEE Trans. on Signal Processing*, vol. 46, no. 12, pp. 3256–3268, Dec. 1998.
- [9] Z. Cvetković and M. Vetterli, “Oversampled filter banks,” *IEEE Trans. Signal Processing*, vol. 46, no. 5, pp. 1245–1255, May 1998.
- [10] P. J. S. G. Ferreira, “Mathematics for multimedia signal processing II — discrete finite frames and signal reconstruction,” in *Signal processing for multimedia*, J. S. Byrnes, Ed., pp. 35–54. IOC Press, 1999.
- [11] A. Aldroubi and K. Gröchenig, “Non-uniform sampling and reconstruction in shift-invariant spaces,” *Siam Rev.*, to appear.
- [12] Y. C. Eldar and A. V. Oppenheim, “Orthogonal and projected orthogonal matched-filter detection,” submitted to *IEEE Trans. Signal Processing*, Jan. 2001.
- [13] Y. C. Eldar and A. M. Chan, “Orthogonal and projected orthogonal multiuser detection,” submitted to *IEEE Trans. Inform. Theory*, May 2001.
- [14] V. K. Goyal, J. Kovačević, and M. Vetterli, “Multiple description transform coding: Robustness to erasures using tight frame expansions,” in *Proc. IEEE Int. Symp. Inform. Theory*, Cambridge, MA, Aug. 1998, p. 408.
- [15] D. Gabor, “Theory of communication,” *J. Inst. Elec. Eng.*, vol. 93, pp. 429–439, Nov. 1946.
- [16] H. G. Feichtinger and T. Strohmer, Eds., *Gabor Analysis: Theory, Algorithms, and Applications*, Birkhäuser, 1998.
- [17] I. Daubechies, *Ten Lectures on Wavelets*, Philadelphia, PA: SIAM, 1992.

- [18] D. Slepian, "Group codes for the Gaussian channel," *Bell Sys. Tech. J.*, pp. 575–602, Apr. 1968.
- [19] G. D. Forney, Jr., "Geometrically uniform codes," *IEEE Trans. Inform. Theory*, vol. 37, pp. 1241–1260, Sep. 1991.
- [20] G. Ungerboeck, "Channel coding with multilevel/phase signals," *IEEE Trans. Inform. Theory*, vol. 28, pp. 55–67, 1982.
- [21] D. Raphaeli, "On multidimensional coded modulations having uniform error property for generalized decoding and flat-fading channels," *IEEE Trans. Commun.*, vol. 46, pp. 34–40, 1998.
- [22] G. Kaiser, *A Friendly Guide to Wavelets*, Boston: Birkhauser, 1994.
- [23] H. Bölcskei and A. J. E. M. Janssen, "Gabor frames, unimodularity, and window decay," *J. Fourier Analys. Appl.*, vol. 6, no. 3, pp. 255–276, 2000.
- [24] A. J. E. M. Janssen and H. Bölcskei, "Equivalence of two methods for constructing tight Gabor frames," *IEEE Signal Processing Lett.*, vol. 7, pp. 79–82, Apr. 2000.
- [25] Y. C. Eldar and G. D. Forney, Jr., "On quantum detection and the square-root measurement," *IEEE Trans. Inform. Theory*, vol. 47, pp. 858–872, Mar. 2001.
- [26] A. J. E. M. Janssen and T. Strohmer, "Characterization and computation of canonical tight windows for Gabor frames," <http://xxx.lanl.gov/abs/math.FA/0010245>.
- [27] Y. Meyer, "Ondelettes et fonctions splines," Dec. 1986, Seminaire EDP, Ecole Polytechnique, Paris, France.
- [28] Y. C. Eldar and G. D. Forney, Jr., "Optimal tight frames and quantum measurement," submitted to *IEEE Trans. Inform. Theory*, June 2001; also available at <http://xxx.lanl.gov/abs/quant-ph/0106070>.
- [29] T. Strohmer, *Finite and infinite-dimensional models for oversampled filter banks*, pp. 297–320, in *Modern Sampling Theory: Mathematics and Applications*, J. J. Benedetto and P. J. S. G. Ferreira, Eds. Boston, MA: Birkhäuser, 2000.
- [30] I. Daubechies, "Orthonormal bases of compactly supported wavelets," *Comm. Pure Appl. Math.*, pp. 909–996, 1988.
- [31] S. G. Mallat, "A theory of multiresolution signal decomposition: The wavelet representation," *IEEE Trans. Pattern Anal. Mach. Intell.*, vol. 11, pp. 674–693, 1989.
- [32] M. Unser and A. Aldroubi, "Families of multiresolution and wavelet spaces with optimal properties," *Numer. Funct. Anal. Optimiz.*, vol. 14, pp. 417–446, 1993.
- [33] A. Shokrollahi, B. Hassibi, B. M. Hochwald, and W. Sweldens, "Representation theory for high-rate multiple-antenna code design," submitted to *IEEE Trans. Inform. Theory*, 2000.

- [34] A. Ben-Israel and T. N. E. Greville, *Generalized Inverses: Theory And Applications*, Huntington, New York: Krieger, 1980.
- [35] I. Daubechies, S. Jaffard, and J. L. Journé, “A simple Wilson orthonormal basis with exponential decay,” *SIAM J. Math. Anal.*, vol. 22, pp. 554–572, 1991.
- [36] M. A. Armstrong, *Groups and Symmetry*, New York: Springer-Verlag, 1988.
- [37] Y. C. Eldar, “Least-squares inner product shaping,” submitted to *Linear Algebra Appl.*, Jan. 2001.
- [38] C. W. Helstrom, *Quantum Detection and Estimation Theory*, New York: Academic Press, 1976.
- [39] A. S. Holevo, “Statistical decisions in quantum theory,” *J. Multivar. Anal.*, vol. 3, pp. 337–394, Dec. 1973.
- [40] H. P. Yuen, R. S. Kennedy, and M. Lax, “Optimum testing of multiple hypotheses in quantum detection theory,” *IEEE Trans. Inform. Theory*, vol. IT-21, pp. 125–134, Mar. 1975.
- [41] M. Charbit, C. Bendjaballah, and C. W. Helstrom, “Cutoff rate for the m -ary PSK modulation channel with optimal quantum detection,” *IEEE Trans. Inform. Theory*, vol. 35, pp. 1131–1133, Sep. 1989.
- [42] M. Ban, K. Kurukow, R. Momose, and O. Hirota, “Optimum measurements for discrimination among symmetric quantum states and parameter estimation,” *Int. J. Theor. Phys.*, vol. 36, pp. 1269–1288, 1997.
- [43] C. W. Helstrom, “Bayes-cost reduction algorithm in quantum hypothesis testing,” *IEEE Trans. Inform. Theory*, vol. 28, pp. 359–366, Mar. 1982.
- [44] G. James and M. Liebeck, *Representations and Characters of Groups*, Cambridge, England: Univ. Press, 1993.
- [45] A. Terras, *Fourier Analysis on Finite Groups and Applications*, Cambridge, UK: Cambridge Univ. Press, 1999.